

Concussions Return to Play

Gradual Return to Sport

When an athlete is returning to sport they should follow a stepwise progression managed by a healthcare provider with increasing amounts of exercise. **Make sure there is at least 24 hours between each step.**

Step 1

Symptom Limited Activity

May do daily activities that do not worsen symptoms.

Goal: Gradual reintroduction of work/school activities.



Step 2

Light Aerobic Exercise

May do walking/stationary bike at slow to medium pace. No resistance training.

Goal: Increased heart rate.



Step 3

Sport Specific Exercise

Running or skating drills. No head impact activities.

Goal: Add movement

Step 4

Non-Contact Training Drills

Harder training drills, i.e., passing drills. May start progressive resistance training (weights).

Goal: Exercise, coordination, and increased thinking.



Step 5

Full Contact Practice

Following medical clearance, participate in normal training activities.

Goal: Restore confidence and assess functional skills by coaching staff.



Return to Play/Sport Normal Game Play

Written Clearance should be provided by a healthcare provider before the athlete begins the return to play protocol as directed by state laws and regulations.

If symptoms get worse during a stage, the athlete should go back to the previous stage until symptom free.



Barrow Concussion and Brain Injury Center 602-406-HEAD (4323)

<https://concussion.barrowneuro.org> | www.barrowneuro.org

Contact: 602-406-4762 | concussiondiscussion@barrowneuro.org

Content source: www.cdc.gov/concussions, McCrory P, Meeuwse W, Dvorak J, et al. Consensus statement on concussion in sport- international conference on concussion in sport held in Berlin, October 2016. Br J Sports Med Published First: 26/4/2017. Doi:10.1136/bjsports-2017-097699