

## Post-Operative Feeding Guidelines Following Lip Repair for Infants with Cleft Lip and Alveolus (Gum Line)

### *How should I feed my infant after surgery?*

Your infant can immediately return to whichever cleft-adapted feeding method you were using prior to the repair. A cleft-adapted bottle/nipple is recommended for one month after surgery. After one month, you can switch your child to a non-adapted bottle/nipple or breastfeeding.

### *How will surgery impact offering purees?*

It is recommended purees be introduced around 6 months of age. If your infant has already started purees prior to lip repair, s/he can immediately return to purees with a spoon. If your infant has not yet started purees prior to the repair, it is recommended to wait to introduce purees until age appropriate, typically 6 months of age.

### *Is my infant allowed to have a pacifier after surgery?*

Pacifiers should not be used for 14 days (2 weeks) after surgery. It is best to wean the infant off of the pacifier prior to surgery. You can try poking a hole in the bottom of pacifier for a few days and then slowly cut the pacifier back every couple of days until the infant can no longer suck on the pacifier and becomes less interested in it.

### *What if my infant is not eating after surgery?*

First, give your infant some time. Sometimes coming out of the anesthesia can be difficult. Start with offering the same bottle that you were using before surgery. If your infant continues to have difficulty, try using one nipple with a larger number (faster flow) on the Dr. Brown Specialty Feeding System or one flow line faster on the Haberman bottle. If your infant continues to have difficulty, request a consultation with a speech-language pathologist.

### *Do I need to be seen back after surgery for feeding therapy?*

Your infant should be seen back at Barrow Cleft and Craniofacial Center one week after surgery and again one month after surgery.

Summary for after surgery:

	<b>Liquids</b>	<b>Purees</b>	<b>Pacifier</b>
<b>Post lip repair</b>	Immediately return to preoperative cleft adapted bottle/nipple Covert to regular nipple/bottle/breast 1 month after surgery	Immediately return to purees with spoon as age appropriate	No pacifier for 14 days

## Barrow Cleft and Craniofacial Center



Jessica Williams, MS, CCC-SLP  
Speech Language Pathologist  
Jessica.Williams4@dignityhealth.org  
602.406.8514

Kelly Nett Cordero, PhD, CCC-SLP  
Bilingual Speech Language Pathologist  
Kelly.Cordero@dignityhealth.org  
602.406.5734